

### 317(6): Self Consistency of Calculations.

The basic geometry of the homogeneous field equation is:

$$D_\mu \tilde{T}^{a\mu\nu} := \tilde{R}_\mu^{a\mu\nu} - (1)$$

which is the Hodge dual form of the Cartan identity.

Therefore: 
$$\partial_\mu \tilde{T}^{a\mu\nu} := \tilde{R}_\mu^{a\mu\nu} - \omega_{\mu b}^a \tilde{T}^{b\mu\nu} - (2)$$

as in Note 255 (6). Here, for each  $a$ :

$$\tilde{T}^{a\mu\nu} := \begin{bmatrix} 0 & -T^1(sp) & -T^2(sp) & -T^3(sp) \\ T^1(sp) & 0 & T^3(orb) & -T^2(orb) \\ T^2(sp) & -T^3(orb) & 0 & T^1(orb) \\ T^3(sp) & T^2(orb) & -T^1(orb) & 0 \end{bmatrix} - (3)$$

and similarly for  $\tilde{R}^a_{b\mu\nu}$ .

### The Gauss Law of Magnetism

In this case: 
$$n = 0, - (4)$$

so: 
$$\partial_1 \tilde{T}^{a10} + \partial_2 \tilde{T}^{a20} + \partial_3 \tilde{T}^{a30}$$

$$:= \sqrt{\frac{1}{2}} \tilde{R}^a_b{}^{10} + \sqrt{\frac{1}{2}} \tilde{R}^a_b{}^{20} + \sqrt{\frac{1}{2}} \tilde{R}^a_b{}^{30} - (5)$$

$$- (\omega^a_{1b} \tilde{T}^{b10} + \omega^a_{2b} \tilde{T}^{b20} + \omega^a_{3b} \tilde{T}^{b30})$$

In eq. (5):

$$d_\mu = \left( \frac{1}{c} \frac{\partial}{\partial t}, \underline{\nabla} \right), \quad - (6)$$

$$\underline{v}_\mu^a = (\underline{v}_0^a, -\underline{v}^a) \quad - (7)$$

$$\underline{\omega}_{\mu b}^a = (\underline{\omega}_{0b}^a, -\underline{\omega}_b^a) \quad - (8)$$

So eq. (5) becomes:

$$\underline{\nabla} \cdot \underline{T}^a(\text{spin}) := \underline{\omega}^a_b \cdot \underline{T}^b(\text{spin}) - \underline{v}^b \cdot \underline{R}^a_b(\text{spin})$$

This is the same as Note 255(6), Q.E.D.<sup>(9)</sup>

Now use:

$$\underline{B}^a = A^{(0)} \underline{T}^a(\text{spin}) \quad - (10)$$

$$\underline{A}^a = A^{(0)} \underline{v}^a \quad - (11) \quad - (12)$$

to obtain:

$$\underline{\nabla} \cdot \underline{B}^a := \underline{\omega}^a_b \cdot \underline{B}^b - \underline{A}^b \cdot \underline{R}^a_b(\text{spin})$$

This is the same as the Engineering Model QED.

Now remove a indices using:

$$\underline{B} = -e_a \underline{B}^a \quad - (13)$$

$$\underline{\omega}_b = -e_a \underline{\omega}^a_b \quad - (14)$$

$$\underline{R}_b = -e_a \underline{R}^a_b \quad - (15)$$

3) and use the E(2) hypothesis:

$$\underline{B}^a_b = W^{(0)} \underline{R}^a_b(\text{spin}) - (16)$$

so:  $\underline{\nabla} \cdot \underline{B}^a = \underline{\omega}^a_b \cdot \underline{B}^b - \frac{1}{W^{(0)}} \underline{A}^b \cdot \underline{B}^a - (17)$

and  $\underline{\nabla} \cdot \underline{B} = \underline{\omega}_b \cdot \underline{B}^b - \frac{1}{W^{(0)}} \underline{A}^b \cdot \underline{B}_b - (18)$

Finally remove the b indices using:

$$\underline{\nabla} \cdot \underline{B} = e_b e^b \underline{\omega} \cdot \underline{B} - \frac{e^b e_b}{W^{(0)}} \underline{A} \cdot \underline{B} - (19)$$

where  $e_b e^b = e^b e_b = -2 - (20)$

so

$$\boxed{\underline{\nabla} \cdot \underline{B} = 2 \left( \frac{1}{W^{(0)}} \underline{A} - \underline{\omega} \right) \cdot \underline{B} - (21)}$$

$$= 2 \left( \frac{v}{r^{(0)}} - \underline{\omega} \right) \cdot \underline{B}$$

There is a sign change for UFT 316 but this does not affect the arguments of UFT 317.

The Faraday law of induction is calculated by way of indices

$$\omega = 1, 2, 3 - (22)$$

For  $n = 1$ :

$$\begin{aligned} & \partial_0 \tilde{T}^{a01} + \partial_2 \tilde{T}^{a21} + \partial_3 \tilde{T}^{a31} \\ &= \sqrt{g}_0^b \tilde{R}^a_b{}^{01} + \sqrt{g}_2^b \tilde{R}^a_b{}^{21} + \sqrt{g}_3^b \tilde{R}^a_b{}^{31} \\ & - \left( \omega^a_{0b} \tilde{T}^{a01} + \omega^a_{2b} \tilde{T}^{b21} + \omega^a_{3b} \tilde{T}^{b31} \right) \end{aligned} \quad (24)$$

i.e.:

$$\begin{aligned} & -\partial_0 T^1(s_p) - \partial_2 T^3(\alpha b) + \partial_3 T^2(\alpha b) \\ &= -\sqrt{g}_0^b R^a_b{}^1(s_p) + \sqrt{g}_2^b R^a_b{}^3(\alpha b) - \sqrt{g}_3^b R^a_b{}^2(\alpha b) \\ & + \omega^a_{0b} T^{a1}(s_p) - \omega^a_{2b} T^{b3}(\alpha b) + \omega^a_{3b} T^{b2}(\alpha b) \end{aligned} \quad (25)$$

This means:

$$\begin{aligned} & -\frac{1}{c} \frac{\partial T_x}{\partial t}(s_p) - \frac{\partial T_z}{\partial y}(\alpha b) + \frac{\partial T_y}{\partial z}(\alpha b) \\ &= -\sqrt{g}_0^b R^a_b{}^x(s_p) - \sqrt{g}_y^b R^a_b{}^z(\alpha b) + \sqrt{g}_z^b R^a_b{}^y(\alpha b) \\ & + \omega^a_{0b} T^a_x(s_p) + \omega^a_{yb} T^b_z(\alpha b) - \omega^a_{zb} T^b_y(\alpha b) \end{aligned} \quad (26)$$

Now use:

$$\underline{A} \times \underline{B} = \begin{vmatrix} \underline{i} & \underline{j} & \underline{k} \\ A_x & A_y & A_z \\ B_x & B_y & B_z \end{vmatrix} \quad (27)$$

5) so:

$$(\underline{A} \times \underline{B})_x = A_y B_z - A_z B_y. \quad (28)$$

It follows that:

$$-\frac{1}{c} \frac{\partial T_x(s_p)}{\partial t} - (\underline{\nabla} \times \underline{T}(arb))_x = -\underline{v}_0^b R^a_{bx}(s_p) - (\underline{v}^b \times \underline{R}^a_b(arb))_x \\ + \omega^a_{ob} T^b_x(s_p) + (\underline{\omega}^a_b \times \underline{T}^b)_x \quad (29)$$

Using similar results for  $y$  and  $z$  components it follows that:

$$\boxed{\begin{aligned} & \frac{1}{c} \frac{\partial \underline{T}^a(s_p)}{\partial t} + \underline{\nabla} \times \underline{T}^a(arb) \\ &= \underline{v}_0^b R^a_b(s_p) + \underline{v}^b \times \underline{R}^a_b(arb) \\ & \quad - \omega^a_{ob} \underline{T}^b(s_p) - \underline{\omega}^a_b \times \underline{T}^b(arb) \end{aligned}} \quad (29)$$

This is the same as eq. (40) of Note 255(1),  
A.E.D.

Now use:  $\underline{B}^a = A^{(0)} \underline{T}^a(s_p) \quad (30)$

$$\underline{E}^a = c A^{(0)} \underline{T}^a(arb) \quad (31)$$

and

$$A_{\mu}^a = A^{(0)} \underline{v}_{\mu}^a - (32)$$

$$\text{So: } (A_0^a, -\underline{A}^a) = A^{(0)} (\underline{v}_0^a, -\underline{v}^a) - (33)$$

$$\text{So } A_0^a = A^{(0)} \underline{v}_0^a - (34)$$

$$\text{and } \underline{A}^a = A^{(0)} \underline{v}^a - (35)$$

It follows that:

$$\frac{1}{c A^{(0)}} \frac{\partial \underline{B}^a}{\partial t} + \frac{1}{c A^{(0)}} \underline{\nabla} \times \underline{E}^a = \frac{A_0^b}{A^{(0)}} \underline{R}_b^a(sp) + \frac{A^b}{A^{(0)}} \times \underline{R}_b^a(orb) - (36)$$

$$- \frac{\omega_{ab}^a}{A^{(0)}} \underline{B}^b - \frac{1}{c A^{(0)}} \underline{\omega}_{ab}^a \times \underline{E}^b - (37)$$

i.e.:

$$\boxed{\frac{\partial \underline{B}^a}{\partial t} + \underline{\nabla} \times \underline{E}^a = c \frac{A_0^b}{A^{(0)}} \underline{R}_b^a(sp) + c \frac{A^b}{A^{(0)}} \times \underline{R}_b^a(orb) - c \omega_{ab}^a \underline{B}^b - \underline{\omega}_{ab}^a \times \underline{E}^b}$$

The Feynberg model on page 26 is

7) written as:

$$\underline{J} = -\omega^a{}_b \underline{B}^b + \underbrace{\omega^a{}_b \times \underline{E}^b}_{\text{dotted terms}} + \underbrace{\underline{\Phi}^b{}_0 \cdot \underline{R}^a{}_b(sp)}_{\text{dotted terms}} \underbrace{c \underline{A}^b \times \underline{R}^a{}_b(al)}_{\text{dotted terms}} \quad (38)$$

So there is a sign error in the engineering model in the dotted terms. In the engineering model the units of  $\omega^a{}_b$  are  $c$  times larger than in eq. (37)

$$\text{and} \quad \underline{\Phi}^b{}_0 = c \omega^a{}_b \quad (39)$$

In eq. (37) we can now use:

$$\underline{B}^a{}_b = \underline{W}^{(0)} \underline{R}^a{}_b(sp) \quad (40)$$

$$\underline{E}^a{}_b = c \underline{W}^{(0)} \underline{R}^a{}_b(al) \quad (41)$$

Therefore:

$$\frac{\partial \underline{B}^a}{\partial t} + \underline{\nabla} \times \underline{E}^a = \frac{c \underline{A}^b}{\underline{W}^{(0)}} \underline{B}^a{}_b + \frac{\underline{A}^b}{\underline{W}^{(0)}} \times \underline{E}^a{}_b - c \omega^a{}_b \underline{B}^b - \omega^a{}_b \times \underline{E}^b \quad (42)$$

Now remove the tangent index using the method given already to find that:

$$8) \frac{\partial \underline{B}}{\partial t} + \underline{\nabla} \times \underline{E} = 2 \left[ c \omega_0 - \frac{c A_0}{W^{(0)}} \right] \underline{B} + \left( \underline{\omega} - \frac{\underline{A}^b}{W^{(0)}} \right) \times \underline{E} \quad - (43)$$

i.e.

$$\boxed{\frac{\partial \underline{B}}{\partial t} + \underline{\nabla} \times \underline{E} = 2 \left[ c \left( \omega_0 - \frac{v_0}{r^{(0)}} \right) \underline{B} + \left( \underline{\omega} - \frac{\underline{v}}{r^{(0)}} \right) \times \underline{E} \right]} \quad - (44)$$

This is the same as in Notes for UFT 316 and UFT 317 QED.

The Complete Field Equations

$$\underline{\nabla} \cdot \underline{B} = 2 \underline{B} \cdot \left( \frac{\underline{v}}{r^{(0)}} - \underline{\omega} \right) \quad - (45)$$

$$\underline{\nabla} \cdot \underline{E} = 2 \underline{E} \cdot \left( \frac{\underline{v}}{r^{(0)}} - \underline{\omega} \right) \quad - (46)$$

$$\frac{\partial \underline{B}}{\partial t} + \underline{\nabla} \times \underline{E} = 2 \left[ c \left( \omega_0 - \frac{v_0}{r^{(0)}} \right) \underline{B} + \left( \underline{\omega} - \frac{\underline{v}}{r^{(0)}} \right) \times \underline{E} \right] \quad - (47)$$

$$\underline{\nabla} \times \underline{B} - \frac{1}{c^2} \frac{\partial \underline{E}}{\partial t} = 2 \left[ \left( \frac{v_0}{r^{(0)}} - \omega_0 \right) \frac{\underline{E}}{c} + \left( \frac{\underline{v}}{r^{(0)}} - \underline{\omega} \right) \times \underline{B} \right] \quad - (48)$$