

The Bianchi identity of differential geometry

M.W. Evans^{1*} and H. Eckardt^{2}**

**Alpha Institute for Advanced Studies (AIAS) (www.aias.us); **Unified Physics Institute of Technology (UPITEC) (www.upitec.org)*

It is shown that the second Bianchi equation used by Einstein and Hilbert is incomplete, so that cosmology based on that equation is also incomplete. A great deal of new information can be obtained by deriving the true second Bianchi identity of differential geometry from first Bianchi identity of Cartan. When this done, it is seen that cosmology based on the Einstein Hilbert field equation is a narrow special case in which the torsion is missing. Using the true Bianchi identity, cosmology can be developed entirely in terms of torsion, in a simpler way, and providing more information.

Keywords: Second Bianchi identity of differential geometry, torsion based cosmology, ECE theory.

1 Introduction

Recently, a generally covariant unified field theory has been developed [1–12] directly on standard differential geometry [13] using the Cartan structure equations and Bianchi equations [14]. This theory has been developed in order to suggest a logical geometrical framework for a unified field theory of natural philosophy. This is of course the philosophy upon which relativity theory is based, and asserts that geometry is the basis of natural philosophy. The theory is known as Einstein–Cartan–Evans (ECE) field theory because it aims to complete the well known work of Einstein and Cartan. In Section 2 it is shown that a new identity of differential geometry can be derived from the first Bianchi identity of differential geometry given by Cartan. It is shown that the true first and second Bianchi identities are related, and that both relate curvature to torsion. The traditional second Bianchi equation as used by Einstein and Hilbert [15] to derive the famous field equation is a narrow special case of the true Bianchi identity. So contemporary cosmology and relativity is also a narrow special case

¹e-mail: EMyrone@aol.com

²e-mail: horsteck@aol.com

of what is possible. This conclusion is illustrated by ECE theory, in which the fundamental fields of physics are unified by the use of torsion to represent the electromagnetic field. It has also been shown that torsion is important [1–12] in a purely gravitational context, and is responsible for example for the formation of spiral galaxies.

In Section 3, a new field equation for cosmology is suggested using the novel concept of Noether forms to represent canonical energy momentum density. In contrast to the traditional Noether tensor of the Einstein Hilbert equation, a symmetric tensor, the Noether forms are anti-symmetric in their last two indices and are made proportional to the torsion form. This procedure greatly simplifies and also extends the field equation of Einstein and Hilbert.

2 Derivation of the true second Bianchi identity

The first Bianchi identity as given by Cartan [13] is:

$$D \wedge T^a = d \wedge T^a + \omega_b^a \wedge T^b := R_b^a \wedge q^b \quad (1)$$

where, in conventional notation, T^a is the torsion form, ω_b^a is the spin connection, R_b^a is the curvature or Riemann form, and q^b is the tetrad form. In tensor notation [1–13], Eq. (1) becomes:

$$\begin{aligned} & \partial_\mu \Gamma_{\nu\rho}^\lambda - \partial_\nu \Gamma_{\mu\rho}^\lambda + \Gamma_{\mu\sigma}^\lambda \Gamma_{\nu\rho}^\sigma - \Gamma_{\nu\sigma}^\lambda \Gamma_{\mu\rho}^\sigma \\ & + \partial_\nu \Gamma_{\rho\mu}^\lambda - \partial_\rho \Gamma_{\nu\mu}^\lambda + \Gamma_{\nu\sigma}^\lambda \Gamma_{\rho\mu}^\sigma - \Gamma_{\rho\sigma}^\lambda \Gamma_{\nu\mu}^\sigma \\ & + \partial_\rho \Gamma_{\mu\nu}^\lambda - \partial_\mu \Gamma_{\rho\nu}^\lambda + \Gamma_{\rho\sigma}^\lambda \Gamma_{\mu\nu}^\sigma - \Gamma_{\mu\sigma}^\lambda \Gamma_{\rho\nu}^\sigma \\ & := R_{\rho\mu\nu}^\lambda + R_{\mu\nu\rho}^\lambda + R_{\nu\rho\mu}^\lambda \end{aligned} \quad (2)$$

where $\Gamma_{\nu\rho}^\lambda$ is the gamma connection, and $R_{\rho\mu\nu}^\lambda$ is the Riemann tensor. Eq. (2) shows that the first Bianchi identity of Cartan is a true identity, it identifies the cyclic sum of three Riemann tensors to a cyclic sum of three definitions of the Riemann tensor. So the right hand side is identically equal to the left hand side, as required of a true identity. Note carefully that Eq. (2) is true for any kind of gamma connection, not just the Christoffel connection. The differential form notation is much more concise and elegant than the tensor notation, but both contain the same information. Implied in both equations (1) and (2) is a non-zero torsion tensor [1–13]:

$$T_{\mu\nu}^\kappa = \Gamma_{\mu\nu}^\kappa - \Gamma_{\nu\mu}^\kappa \neq 0. \quad (3)$$

This vanishes for the symmetric Christoffel connection:

$$\Gamma_{\mu\nu}^{\kappa} = \Gamma_{\nu\mu}^{\kappa}. \quad (4)$$

The first Bianchi equation used by Einstein and Hilbert, and in conventional cosmology, is

$$R_b^a \wedge q^b = 0 \quad (5)$$

and this is not an identity because it is true if and only if the metric is symmetric, i.e. for a Christoffel connection. Eq. (5) is often mistakenly known as "the first Bianchi identity", but it is not a true identity. It was actually discovered [13] by Ricci and Levi-Civita, and not by Bianchi. In tensor notation, Eq. (5) is:

$$R_{\rho\sigma\mu\nu} + R_{\rho\mu\nu\sigma} + R_{\rho\nu\sigma\mu} = 0 \quad (6)$$

where:

$$R_{\rho\sigma\mu\nu} = g_{\rho\kappa} R_{\sigma\mu\nu}^{\kappa} \quad (7)$$

is the Riemann tensor with indices lowered. Here

$$g_{\rho\kappa} = g_{\kappa\rho} \quad (8)$$

is the symmetric metric. Note that Eqs (5) and (6) imply a vanishing torsion:

$$T_{\mu\nu}^{\kappa} = \Gamma_{\mu\nu}^{\kappa} - \Gamma_{\nu\mu}^{\kappa} = 0. \quad (9)$$

In Riemann normal coordinates [13] Eq. (6) is:

$$\begin{aligned} & R_{\alpha\beta\gamma\delta} + R_{\alpha\gamma\delta\beta} + R_{\alpha\gamma\beta\delta} \\ &= \frac{1}{2} \left(\partial_{\rho} \partial_{\gamma} g_{\alpha\delta} - \partial_{\alpha} \partial_{\gamma} g_{\beta\delta} - \partial_{\beta} \partial_{\delta} g_{\alpha\gamma} + \partial_{\alpha} \partial_{\gamma} g_{\beta\gamma} \right) \\ &\quad + \partial_{\gamma} \partial_{\delta} g_{\alpha\beta} - \partial_{\alpha} \partial_{\delta} g_{\gamma\beta} - \partial_{\gamma} \partial_{\beta} g_{\alpha\delta} + \partial_{\alpha} \partial_{\beta} g_{\gamma\delta} \\ &\quad + \partial_{\delta} \partial_{\beta} g_{\alpha\gamma} - \partial_{\alpha} \partial_{\beta} g_{\alpha\gamma} - \partial_{\alpha} \partial_{\gamma} g_{\alpha\beta} + \partial_{\alpha} \partial_{\gamma} g_{\delta\beta} \\ &= 0 \end{aligned} \quad (10)$$

and it is seen that this is not true of the metric is not symmetric, and is not true if the connection is not a Christoffel connection. For the general connection the true first Bianchi identity is Eq. (1) or Eq. (2) and Eq. (1) is the field equation

of classical electrodynamics in ECE theory [1–12].

A second Bianchi equation of Cartan geometry is given in ref. [13] as:

$$D \wedge R_b^a = 0 \quad (11)$$

but this is a special case [13] when the torsion is zero (the Einstein Hilbert case). In tensor notation Eq. (11) is:

$$D_\lambda R_{\rho\sigma\mu\nu} + D_\rho R_{\sigma\lambda\mu\nu} + D_\sigma R_{\lambda\rho\mu\nu} = 0 \quad (12)$$

and is always referred to as "the second Bianchi identity" in conventional cosmology. In fact it is not an identity. This fact is shown [13] by expanding it in Riemann normal coordinates:

$$\begin{aligned} & D_\lambda R_{\rho\sigma\mu\nu} + D_\rho R_{\sigma\lambda\mu\nu} + D_\sigma R_{\lambda\rho\mu\nu} \\ &= \frac{1}{2} \left(\partial_\lambda \partial_\mu \partial_{\sigma\rho\nu} - \partial_\lambda \partial_\mu \partial_{\rho\sigma\nu} - \partial_\lambda \partial_\nu \partial_{\sigma\rho\mu} - \partial_\lambda \partial_\nu \partial_{\rho\sigma\mu} \right. \\ & \quad + \partial_\rho \partial_\mu \partial_{\lambda\sigma\nu} - \partial_\rho \partial_\mu \partial_{\sigma\nu\lambda} - \partial_\rho \partial_\nu \partial_{\lambda\sigma\mu} + \partial_\rho \partial_\nu \partial_{\sigma\mu\lambda} \\ & \quad \left. + \partial_\sigma \partial_\mu \partial_{\rho\sigma\lambda\nu} - \partial_\sigma \partial_\mu \partial_{\lambda\sigma\nu\rho} - \partial_\sigma \partial_\nu \partial_{\rho\sigma\lambda\mu} + \partial_\sigma \partial_\nu \partial_{\lambda\sigma\mu\rho} \right) \\ &= 0. \end{aligned} \quad (13)$$

It is seen by comparison of the dotted terms in the above equation that a zero result is obtained if and only if:

$$g_{\rho\nu} = g_{\nu\rho} \quad (14)$$

where the commutation of partial four-derivatives has been used:

$$\partial_\lambda \partial_\mu \partial_\sigma = \partial_\mu \partial_\sigma \partial_\lambda, \quad (15)$$

$$\partial_\mu \partial_\sigma \partial_\lambda = \partial_\sigma \partial_\mu \partial_\lambda. \quad (16)$$

Eq. (14) is true if and only if the connection is the Christoffel connection:

$$\Gamma_{\mu\nu}^\kappa = \Gamma_{\nu\mu}^\kappa \quad (17)$$

and if and only if the torsion tensor vanishes:

$$T_{\mu\nu}^\kappa = \Gamma_{\mu\nu}^\kappa - \Gamma_{\nu\mu}^\kappa = 0. \quad (18)$$

Equation (12) can be rewritten using contraction of indices [13] as:

$$D^\mu G_{\nu\mu} = 0 \tag{19}$$

where $G_{\nu\mu}$ is the Einstein tensor:

$$G_{\nu\mu} = R_{\nu\mu} - \frac{1}{2}Rg_{\nu\mu}. \tag{20}$$

Here $R_{\nu\mu}$ is the symmetric Ricci tensor [13] and R is the conventional scalar curvature of cosmology [13]. The Einstein Hilbert field equation is obtained by making Eq. (19) proportional to the Noether Theorem:

$$D^\mu T_{\nu\mu} = 0 \tag{21}$$

i.e.

$$D^\mu G_{\nu\mu} = kD^\mu T_{\nu\mu} \tag{22}$$

and using the solution:

$$G_{\nu\mu} = kT_{\nu\mu} \tag{23}$$

which is the Einstein Hilbert field equation of 1915. Here:

$$T_{\nu\mu} = T_{\mu\nu} \tag{24}$$

is the symmetric canonical energy momentum tensor of Noether, and Eq. (21) denotes conservation of energy – momentum as is well known.

It is seen that the famous field equation and the equally famous "second Bianchi identity" are true if and only if the connection is symmetric and if and only if the torsion tensor is zero. They are no longer true otherwise, so conventional cosmology is severely constrained by these assumptions [1–12].

The true second Bianchi identity is obtained by taking the $D \wedge$ derivative of both sides of the true first Bianchi identity (1). Thus:

$$D \wedge (R_b^a \wedge q^b) := D \wedge (D \wedge T^a). \tag{25}$$

The general rule for the exterior derivative of an n -form is [13]

$$(d \wedge A)_{\mu_1 \dots \mu_{\rho+1}} = (\rho + 1) \partial_{[\mu_1 \wedge \mu_2 \dots \mu_{\rho+1}]} \quad (26)$$

where the [] denote anti-symmetric permutation [13] defined as follows:

$$T_{[\mu_1 \mu_2 \dots \mu_n]} = \frac{1}{n!} (T_{\mu_1 \mu_2 \dots \mu_n} + \text{antisymmetrized sum}). \quad (27)$$

Permutations which are the result of an odd number of exchanges are given a minus sign. Thus:

$$T_{[\mu\nu]} = \frac{1}{2} (T_{\mu\nu} - T_{\nu\mu}) \quad (28)$$

and:

$$T_{[\mu\nu\rho]} = \frac{1}{6} (T_{\mu\nu\rho} - T_{\mu\rho\nu} + T_{\rho\mu\nu} - T_{\nu\mu\rho} + T_{\nu\rho\mu} - T_{\rho\nu\mu}). \quad (29)$$

If we assume anti-symmetry in the last two indices:

$$T_{\mu\nu\rho} = -T_{\mu\rho\nu} \text{ etc.} \quad (30)$$

then Eq. (29) becomes:

$$T_{[\mu\nu\rho]} = \frac{1}{3} (T_{\mu\nu\rho} + T_{\rho\mu\nu} + T_{\nu\rho\mu}). \quad (31)$$

When there are four indices the permutation rule is the same as for the subscripts of the rank four Levi-Civita tensor:

$$\begin{aligned} 0123 &= 0312 = 0231 = 1 \\ 1203 &= 1320 = 1032 = 1 \\ 2301 &= 2013 = 2130 = 1 \\ 3102 &= 3210 = 3021 = 1 \\ 0321 &= 0213 = 0132 = -1 \\ 1302 &= 1023 = 1230 = -1 \\ 2103 &= 2310 = 2031 = -1 \\ 3201 &= 3012 = 3120 = -1. \end{aligned} \quad (32)$$

So with:

$$\sigma = 0, \mu = 1, \nu = 2, \rho = 3 \quad (33)$$

and assuming anti-symmetry in the last two indices we obtain:

$$\begin{aligned} T_{[\sigma\mu\nu\rho]} = & \frac{1}{12} (T_{\sigma\mu\nu\rho} + T_{\sigma\rho\mu\nu} + T_{\sigma\nu\rho\mu} \\ & + T_{\mu\nu\sigma\rho} + T_{\mu\rho\nu\sigma} + T_{\mu\sigma\rho\nu} \\ & + T_{\nu\rho\sigma\mu} + T_{\nu\sigma\rho\mu} + T_{\nu\mu\rho\sigma} \\ & + T_{\rho\sigma\mu\nu} + T_{\rho\nu\mu\sigma} + T_{\rho\sigma\nu\mu}) \end{aligned} \quad (34)$$

Thus:

$$\begin{aligned} (D \wedge R)_{\text{cyclic}} & := D \wedge (R^a \wedge q^b) = D \wedge R^a_{\mu\nu\rho} + D \wedge R^a_{\nu\rho\mu} + D \wedge R^a_{\rho\mu\nu} \\ & \frac{1}{12} \left((D_\sigma R^a_{\mu\nu\rho} + D_\sigma R^a_{\rho\mu\nu} + D_\sigma R^a_{\nu\rho\mu} + D_\mu R^a_{\nu\sigma\rho} + D_\mu R^a_{\rho\nu\sigma} + D_\mu R^a_{\sigma\rho\nu}) \right. \\ & + D_\rho R^a_{\sigma\mu\nu} + D_\rho R^a_{\nu\sigma\mu} + D_\rho R^a_{\mu\nu\sigma} + D_\nu R^a_{\rho\sigma\mu} + D_\nu R^a_{\sigma\mu\rho} + D_\nu R^a_{\mu\rho\sigma}) \\ & + (D_\sigma R^a_{\nu\rho\mu} + D_\sigma R^a_{\mu\nu\rho} + D_\sigma R^a_{\rho\mu\nu} + D_\nu R^a_{\rho\sigma\mu} + D_\nu R^a_{\mu\rho\sigma} + D_\nu R^a_{\sigma\mu\rho}) \\ & + D_\rho R^a_{\mu\sigma\nu} + D_\rho R^a_{\sigma\nu\mu} + D_\rho R^a_{\nu\mu\sigma} + D_\mu R^a_{\sigma\nu\rho} + D_\mu R^a_{\rho\sigma\nu} + D_\mu R^a_{\nu\rho\sigma}) \\ & + (D_\sigma R^a_{\rho\mu\nu} + D_\sigma R^a_{\nu\rho\mu} + D_\sigma R^a_{\mu\nu\rho} + D_\rho R^a_{\mu\sigma\nu} + D_\rho R^a_{\nu\mu\sigma} + D_\rho R^a_{\sigma\nu\mu}) \\ & \left. + D_\mu R^a_{\nu\sigma\rho} + D_\mu R^a_{\rho\nu\sigma} + D_\mu R^a_{\sigma\rho\nu} + D_\nu R^a_{\sigma\rho\mu} + D_\nu R^a_{\mu\sigma\rho} + D_\nu R^a_{\rho\mu\sigma}) \right) \\ & = \frac{1}{4} (D_\sigma R^a_{\mu\nu\rho} + D_\sigma R^a_{\rho\mu\nu} + D_\sigma R^a_{\nu\rho\mu}) \\ & + \frac{1}{12} (D_\nu R^a_{\rho\sigma\mu} + D_\nu R^a_{\mu\rho\sigma} + D_\nu R^a_{\sigma\mu\rho} \\ & + D_\rho R^a_{\mu\sigma\nu} + D_\rho R^a_{\sigma\nu\mu} + D_\rho R^a_{\nu\mu\sigma} \\ & + D_\mu R^a_{\nu\sigma\rho} + D_\mu R^a_{\rho\nu\sigma} + D_\mu R^a_{\sigma\rho\nu}) \end{aligned} \quad (35)$$

which is an antisymmetrized sum of terms which we denote as $(D \wedge R)_{\text{cyclic}}$ in condensed notation defined by Eq. (35). Similarly:

$$\begin{aligned}
(D(D \wedge T))_{\text{cyclic}} &:= D \wedge (D \wedge T^a) \\
&= \frac{1}{4} (D_\sigma D_\mu T_{\nu\rho}^a + D_\sigma D_\rho T_{\mu\nu}^a + D_\sigma D_\nu T_{\rho\mu}^a) \\
&\quad + \frac{1}{12} (D_\nu D_\rho T_{\sigma\mu}^a + D_\nu D_\mu T_{\rho\sigma}^a + D_\nu D_\sigma T_{\mu\rho}^a \\
&\quad + D_\rho D_\mu T_{\sigma\nu}^a + D_\rho D_\sigma T_{\nu\mu}^a + D_\rho D_\nu T_{\mu\sigma}^a \\
&\quad + D_\mu D_\nu T_{\sigma\rho}^a + D_\mu D_\rho T_{\nu\sigma}^a + D_\mu D_\sigma T_{\rho\nu}^a)
\end{aligned} \tag{36}$$

The overall result is non-trivial, and is the true Bianchi identity. This Theorem may be stated as follows. If:

$$D \wedge T^a := R_b^a \wedge q^b \tag{37}$$

it follows identically that:

$$(D \wedge R)_{\text{cyclic}} := (D(D \wedge T))_{\text{cyclic}}. \tag{38}$$

Equations (37) and (38) denote the true second Bianchi identity. So the complete set of Cartan equations are:

$$\begin{aligned}
T &= D \wedge q, \\
R &= D \wedge \omega, \\
D \wedge T &:= R \wedge q, \\
(D \wedge R)_{\text{cyclic}} &:= (D(D \wedge T))_{\text{cyclic}}.
\end{aligned} \tag{39}$$

The Einstein–Hilbert field theory is the special case:

$$R \wedge q = 0 \tag{40}$$

$$D \wedge R = 0 \tag{41}$$

and misses a great deal of basic information. It is seen that there is only one true Bianchi identity because Eq. (38) has been derived from Eq. (37).

3 Torsion based cosmology

From the foregoing it is now known that there is only one true Bianchi identity:

$$D \wedge T^a := R_b^a \wedge q^b \tag{42}$$

because this implies the identity:

$$(D \wedge R)_{\text{cyclic}} := (D(D \wedge T))_{\text{cyclic}} \tag{43}$$

In deriving his 1915 field equation, Einstein used the special case:

$$R_b^a \wedge q^b = 0. \tag{44}$$

Using Eq. (37), i.e.:

$$D \wedge (R_b^a \wedge q^b) = 0 \tag{45}$$

it is seen that the special case (44) implies:

$$D \wedge R_b^a = 0 \tag{46}$$

which is the conventional "second Bianchi identity". It has been derived from "the first Bianchi identity" (44) and so the two identities are not independent. The Einstein field equation of 1915 is, in differential form notation:

$$D \wedge R_b^a = kD \wedge N_b^a \tag{47}$$

where k is the Einstein constant, and where we define N_b^a is the Noether form. The latter is a tensor valued two-form representing canonical energy momentum density. The particular solution chosen by Einstein in 1915 is equivalent in form notation to:

$$R_b^a = kN_b^a. \tag{48}$$

There are two Noether forms, related by the structure of Eq. (42), the true Bianchi identity. Thus:

$$D \wedge N^a := N_b^a \wedge q^b \tag{49}$$

where:

$$T^a = kN^a. \tag{50}$$

Here:

$$k = \frac{8\pi G}{c^2} = 1.86595 \times 10^{-26} \text{ m kg m}^{-1} \quad (51)$$

where G is Newton's constant. Thus N^a has the units of mass per unit volume (density) and N^a has the units of mass per unit area. Eq. (49) is the generalization of the Noether Theorem to include rotational as well as translational canonical energy momentum density. In the special case:

$$N_b^a \wedge q^b = 0 \quad (52)$$

then:

$$D \wedge N_b^a = 0. \quad (53)$$

Eq. (53) is equivalent [1–13] in tensor notation to:

$$D^\mu N_{\nu\mu} = 0 \quad (54)$$

where $N_{\nu\mu}$ is the well known symmetric canonical energy momentum density tensor of Noether:

$$N_{\nu\mu} = N_{\mu\nu}. \quad (55)$$

Since TOTAL energy momentum is conserved, Eq. (49) describes the conservation and interaction of rotational and translational energy – momentum. The well known Eq. (54) is true only in the absence of rotational energy–momentum and is again a special case. In general both N_b^a and N^a are roto-translational in nature. The most general field equations of the Einstein type are Eqs. (48) and (50), in which there can be interaction between torsion (rotation) and curvature (translation). These considerations are missing completely from the Einstein field equation and standard cosmology, most of whose contemporary conclusion are based in the very restricted Eq. (53) and are therefore at best incomplete, at worst erroneous. Ideas such as Big Bang and dark matter are erroneous for several reasons, notably the complete neglect of torsion in standard cosmology, and also the erroneous treatment of singularities as shown by Crothers [1–12].

The true Noether theorem (49) can be developed as the generally covariant equation [1–13]:

$$d \wedge N^a = j^a = N_b^a \wedge q^b - \omega_b^a \wedge N^b. \quad (56)$$

When there is no torsional energy momentum density present:

$$N^a = 0 \tag{57}$$

so Eq. (56) is:

$$N^a_b \wedge q^b = 0. \tag{58}$$

When the motion is purely torsional (rotational) the following relation holds, a relation that is akin to ECE field theory [1–12]:

$$N^a_b \wedge q^b = \omega^a_b \wedge N^b, \tag{59}$$

In this case N^a_b is the dual of N^a :

$$N^a_b = -\frac{1}{2} \kappa \epsilon^a_{bc} N^c \tag{60}$$

where κ has the units of inverse meters, or wave-number. So for pure rotational or torsional motion N^a_b and N^a are respectively tensor and vector valued two-forms that denote rotational energy momentum densities. In this case the conservation of energy – momentum is given by:

$$d \wedge N^a = 0. \tag{61}$$

This equation can be developed in vector notation as two equations:

$$\nabla \cdot N^a (\text{spin}) = 0, \tag{62}$$

$$\nabla \times N^a (\text{orbital}) + \frac{1}{c} \frac{\partial N^a}{\partial t} (\text{spin}) = 0. \tag{63}$$

The first is akin to the Gauss law of magnetism and the second to the Faraday law of induction. These equations relate orbital (N^a) and spin (N^a) canonical energy – momentum densities. So this type of torsional cosmology, as seen for example in a spiral galaxy [1–12] can be developed entirely without the Riemann tensor, thus simplifying the solutions dramatically.

References

- [1] M.W.Evans, "Generally Covariant Unified Field Theory" (Abramis Academic, Suffolk, 2005), vol. 1.
- [2] M. W. Evans, *ibid.*, vols. 2 and 3 (2006).
- [3] M. W. Evans, *ibid.*, vol. 4 (2007), papers 55 to 70 on www.aias.us.
- [4] M. W. Evans, *ibid.*, vol. 5 (2008), papers 71 to 89 on www.aias.us.
- [5] M. W. Evans, *ibid.*, vol. 6 (2009), in prep.
- [6] L. Felker, "The Evans Equations of Unified Field Theory" (Abramis Academic, Suffolk, 2007).
- [7] H. Eckardt, L. Felker, S. Crothers, D. Indranu, K. Pendergast and G. J. Evans, articles on www.aias.us, F. Amador, article on www.aias.us in prep., also Omnia Opera of this website, with hyperlinks.
- [8] M. W. Evans, *Acta Phys. Polonica B*, **38**, 2211 (2007); M. W. Evans and H. Eckardt, *Physica B*, in press (2007).
- [9] M. W. Evans (ed.), "Modern Non-linear Optics", a special topical issue in three parts of I. Prigogine and S. A. Rice, "Advances in Chemical Physics" (Wiley Interscience, New York, 2001), vols. 119(1) to 119(3) endorsed by the Royal Swedish Academy; M. W. Evans and S. Kielich (eds.), *ibid.*, vols. 85(1) to 85(3) (Wiley Interscience, New York, 1992, reprinted 1993 and 1997), Polish Govt. Award for excellence.
- [10] M. W. Evans and L. B. Crowell, "Classical and Quantum Electrodynamics and the B(3) Field" (World Scientific, 2001).
- [11] M. W. Evans et al, *Found. Phys. and Found. Phys. Lett.*, 1994 to present, see Omnia Opera of www.aias.us for a complete list with some hyperlinks.
- [12] M. W. Evans and J.-P. Vigi er, "The Enigmatic Photon" (Kluwer, Dordrecht, 1994 to 2002, hardback and softback) in five volumes.
- [13] S. P. Carroll, "Spacetime and Geometry : an Introduction to General Relativity" (Addison-Wesley, New York, 2004, also 1997 notes on web).
- [14] L. H. Ryder, "Quantum Field Theory" (Cambridge Univ. Press, 1996, 2nd ed.).
- [15] A. Einstein, "The Meaning of Relativity" (Princeton Univ. Press, 1921–1954 Eds.).